

# ANTIBIOTIC TREATMENT VOUCHER (STI)

MUST BE ACCOMPANIED BY A VALID, SIGNED PRESCRIPTION

In the state of Michigan, expedited partner therapy (EPT) is limited to azithromycin and /or cefixime

PLEASE PRESENT  
THIS VOUCHER TO ONE  
OF THE FOLLOWING  
PHARMACIES:

**Benson Drug Sav-Mor**  
961 Spring Street  
Muskegon, MI 49442  
231.722.2861

**Community Care Pharmacy**  
2700 Baker Street  
Muskegon Heights, MI 49444  
231.737.9510

**Mercy Health Pharmacy  
Glenside**  
1663 W. Sherman Boulevard  
Muskegon, MI 49441  
231.755.2443

**Mercy Health Pharmacy  
Hackley**  
1675 Leahy Street  
Suite 111  
Muskegon, MI 49442  
231.728.5888

**Mercy Health Pharmacy  
Lakes Village**  
6401 Prairie Street  
Suite 1100  
Muskegon, MI 49444  
231.727.7968

**Mercy Health Pharmacy  
Westshore**  
1150 E. Sherman Boulevard  
Suite 1400  
Muskegon, MI 49444  
231.672.2204

**Meijer Pharmacies, all  
West Michigan locations**

**Muskegon Family Care  
Pharmacy**  
2201 S. Getty Street  
Muskegon Heights, MI 49444  
231.737.1706

## PATIENT INFORMATION

Patient Name: \_\_\_\_\_

Date of Birth: \_\_\_\_\_ Date: \_\_\_\_\_

Address: \_\_\_\_\_

Phone Number: \_\_\_\_\_

Prescribing Provider (print): \_\_\_\_\_

Provider Location: \_\_\_\_\_

Provider Phone: \_\_\_\_\_

## PRESCRIPTION

Prescriber, CHECK ALL THAT APPLY, including diagnosis and treatment prescribed.

**GONOCOCCAL**

Give Ceftriaxone<sup>1</sup> 250 mg I.M. x 1 dose in the office.

**OR**  Cefixime 400 mg p.o. STAT, **ONLY<sup>2</sup>** if Ceftriaxone is not available

**PLUS**  Azithromycin 1 gm p.o. STAT

If severe penicillin or cephalosporin allergy and patient is not a candidate for I.M. Ceftriaxone or p.o. Cefixime consider Gentamicin<sup>1</sup> 240mg I.M. x1 dose in the office.

**AND**  Azithromycin **2 gm** p.o. STAT

- Injectable medications are not paid for by this voucher program and should be billed to the patient/their insurance as per established office policy and procedure.*
- Cefixime should only be considered an alternative to ceftriaxone because it does not provide as high, nor sustained bactericidal blood levels as a 250 mg dose of ceftriaxone and demonstrates limited efficacy for pharyngeal gonorrhea. Any person with pharyngeal gonorrhea who is treated with cefixime should return 14 days after treatment for a test-of cure.*

**C. TRACHOMATIS** (Note: Ceftriaxone is not indicated for treatment of *C. trachomatis* alone)

**Recommended Regimens:**

- Azithromycin 1 gm p.o. STAT, **or**  
 Doxycycline\* 100 mg p.o. BID x 7 day's

\* contraindicated in pregnancy

**Regimen in Pregnancy:**

- Chlamydia recommended regimen:  
 Azithromycin 1 gm p.o. STAT

**Alternative Regimens:**

- Erythromycin 500 mg p.o. QID x 7 days  
 Erythromycin (EES) 800 mg QID x 7 days  
 Levaquin 500 mg p.o. daily x 7 days

**Alternative Regimens for Pregnancy:**

- Amoxicillin 500 mg p.o. TID x 7 days  
 Erythromycin 500 mg p.o. QID x 7 days  
 Erythromycin 250 mg p.o. QID x 14 days  
 Erythromycin (EES) 800 mg QID 7 days  
 Erythromycin (EES) 400 QID x 14 days

No Expiration Date